

Mrk 421: *EUVE* observations from 1994 to 1997

I. Cagnoni

International School for Advanced Studies (SISSA-ISAS), Trieste, Italy

A. Fruscione

Harvard-Smithsonian Center for Astrophysics, Cambridge, MA, USA

I. E. Papadakis

University of Crete, Greece

1. Observations: variability and spectra

From 1994 to 1997 Mrk 421 was observed by *EUVE* (0.02–0.21 keV) 4 times with the Deep Survey/Spectrograph, which allows simultaneous imaging and spectroscopy (e.g. Welsh et al. 1990) and 2 times with the scanner (imaging) telescopes for a total of ~ 1000 ks (see Table 1 for details).

Variability: Not only *EUVE* lightcurves taken from 1994 to 1997 play a key role in multiwavelengths studies, but, since they represent the longest continuous coverages for this source at these energies, they allow a detailed variability analysis. We analyzed all 5 lightcurves in an homogeneous way and we studied the variability of Mrk 421 through the power spectrum technique. We fitted the power spectra with a power law model and the best fit slope found using all the DS lightcurves binned over 5544 s is -2.52 ± 0.40 , while combining the DS and scanner lightcurves we obtain -2.18 ± 0.30 (further details can be found in Cagnoni et al. 1998).

Energy spectra: We analyzed the three spectral data sets in an homogeneous way using the proper off-axis calibrations (Marshall 1998, private communication). We modeled the EUV spectrum of Mrk 421 over the range where the ISM does not absorb all the photons (70–100 Å; 75–110 Å for 1994 observation) with an absorbed power law whose best fit values are given in Table 1. In order to check for the presence and depth of the absorption feature claimed by Kartje et al. 1997 and in an attempt to follow the development of the flare, we divided the 1995 observation into three parts (see Table 1): April 25 - 28 (the flare), April 28 - May 6 (where the absorption feature was claimed) and May 7 - 13.

2. Discussion

Unfortunately the large errors on the slope and normalization make it impossible to follow the spectral development of the flare and even if both the correction for the proper off-axis effective area and the flattening of Mrk 421 spectrum below 1.5 keV of ~ 0.5 (Takahashi private communication) combine to weaken the evidence of the feature around ~ 70 Å, nothing conclusive can be said. The total

Table 1. *EUVE* Observations of Mrk 421 from 1994 to 1997

Observations				Lightcurves		Energy Spectra	
Year	Instr.	Begin	End	T _{exp} (ks)	Count rate (10 ⁻¹ c s ⁻¹)	Slope	Normalization (μJy at 80Å)
94	DS/S	Apr. 2	Apr. 12	280 ^a	1.758 ± 0.009	1.0 ^{+0.6} _{-0.9} ^b	340 ± 18
95	Scanner	Feb. 4	Feb. 7	68	3.219 ± 0.029 ^c	—	—
95	DS/S	Apr. 25	May 13	355	2.914 ± 0.010	1.1 ± 0.4	355 ± 12
95(p. 1)	DS/S	Apr. 25	Apr. 28	86 ^d	3.743 ± 0.023	1.4 ± 0.6	540 ⁺³⁵ ₋₂₀
95(p. 2)	DS/S	Apr. 28	May 6	154 ^d	2.950 ± 0.015	2.2 ± 0.5 ^e	385 ⁺¹⁶ ₋₁₉
95(p. 3)	DS/S	May 7	May 13	115	2.240 ± 0.016	-0.6 ^{+1.0} _{-1.2}	185 ⁺²¹ ₋₁₇
96	DS/S	Apr. 17	Apr. 30	299	3.040 ± 0.011	2.1 ± 0.3	495 ± 15
96	DS/S	May 10	May 11	3.6	4.066 ± 0.113	—	—
97	Scanner	Feb. 7	Feb. 11	108	1.920 ± 0.016 ^c	—	—
Total				1114			

^{a,b}Lightcurve presented in Fruscione et al. 1996; the best fit of the spectral slope was 1.4 ± 0.8

^cCount rate in the Scanner; the equivalent in the DS/S would be 4.506 ± 0.041 in 1995 and 2.690 ± 0.022 in 1997

^{d,e}Lightcurve presented in Kartje et al. 1997; the best fit of the spectral slope was 3.5 ± 0.8

EUVE light curve seems to be smoothly varying on the long time-scale while on a shorter time-scale there is evidence of an *EUVE* flare correlated to the 1995 TeV flare. The best fit value for the slope of the power spectrum is consistent with results found for PKS 2155-304 in the X-rays ($\alpha = -2.5 \pm 0.2$ Tagliaferri et al 1991) and in optical ($\alpha = -2.4$ Paltani et al 1997); it is also consistent with the $\alpha = -2$ expected from a superposition of exponentially decaying/rising shots. A comparison with the jet emission inhomogeneous models (e.g. Celotti et al. 1991) suggests that the shots could be due to perturbations of growing thickness propagating outward along the jet. These perturbations act as an increase of the relativistic electron density and magnetic field strength in the emitting region. Since we do not see any break in the power spectrum, we can place only a lower limit on the decaying time scales of the shots which are likely responsible for the source variability. The limit implies a long decaying time scale ($> 1/10^{-5.5}$ Hz $\simeq 3.66$ days) spread over a narrow range.

Acknowledgments. This work was supported by AXAF S.C. NASA Contract NAS 8-39073 and by NASA grant NAG 5-3174 and NAG 5-3191.

References

- Cagnoni, I., Fruscione, A. & Papadakis, I., 1998, in preparation
 Celotti, A., Maraschi, L. & Treves, A., 1991, ApJ, 377, 404
 Fruscione, A. et al., 1996, in UV and X-ray Spectroscopy of Astroph. and Lab. Plasmas, ed. Yamashita & Watanabe (Tokyo: Univ. Acad. Press), 351
 Kartje J.F., Konigl, A., Hwang, C.Y. & Bowyer, S., 1997, ApJ, 474, 630
 Paltani, S., Courvoisier, T.J.L., Blecha, A. & Bratschi, P., 1997, ApJ, 327, 539
 Tagliaferri, G. et al. 1991, ApJ, 380, 78
 Welsh, B.Y. et al., 1990, Opt. Eng., 29, 752